

Predicting the Response of Insect Pests and Diseases of Arabica Coffee to Climate Change along an Altitudinal Gradient in Mt. Elgon Region, Uganda

Kagezi, H. Godfrey^{1*}, Kucel Patrick¹, Kobusinge Judith¹, Olango D. Nicholas¹
Nakibuule Lilian¹ & Wagoire, W. William¹.

Abstract

Recent studies have predicted an increase in temperature due to climate change which is likely to upsurge pressure from pests and diseases on Arabica coffee, *Coffea arabica*. Under the future climate change scenarios, elevation gradients can be used as analogues for global warming. Basing on this, we therefore determined the incidence and damage of Arabica coffee pests and diseases along an altitudinal gradient in Mt. Elgon region in order to predict their possible impact under these scenarios. Seven insect pest species were recorded - with the white coffee stem borers (WCSB) having the highest incidence of 13%. On the other hand, three diseases were recorded - with coffee leaf rust (CLR) having the highest mean severity of 1.9 on a 1-5 scale. Multiple regression analysis showed that the incidence of only WCSB decreased significantly ($p=0.0169$) with increasing altitude. This implies that under the future climate change scenarios, the impact of this insect pest on Arabica coffee will increase with increase in temperature. Therefore, adaptation mechanisms for climate adversaries such as inter-planting with shade trees should be advanced. However, these tree species should not compete with the coffee as well as promoting pests and diseases.

Key words: Altitudinal-gradient, diseases, climate-change, pests, shade-trees

1. Introduction

Coffee is currently the most important cash crop of Uganda (Musoli et al., 2001). The crop is grown by over 1.3 million households (a quarter of which are headed by females), who are spread out over several production zones (Hill, 2005). There are two main types of coffee grown by farmers in the country, Robusta coffee, *Coffea canephora* Pierre ex A. Foreigner which accounts for about 80% of the coffee and Arabica coffee, *Coffea arabica*, which occupies the remaining 20%. Robusta coffee is predominantly grown below 1200 meters above sea level (m.a.s.l), while Arabica dominates highland regions above 1500 m.a.s.l, particularly in the Mt. Elgon region located in the eastern part of the country (Musoli et al., 2001).

However, production of Arabica coffee in this region is constrained by a number of factors, with pests and diseases are paramount (Musoli et al., 2001; Matovu et al., 2013; Kucel et al., 2016; Liebig et al., 2016). The incidence and severity of these insect pests and diseases differ along the altitudinal gradient (Matovu et al., 2013; Kucel et al., 2016), due to variation in temperature and precipitation. Increase in altitude results into lower temperatures and higher precipitation (Ohmura, 2012). This in turn influences species richness and abundance of insect species as well as plant pathogens, and thus, the damage they cause to plants (Hodkinson, 2005; Deutsch et al., 2008; Matovu et al., 2013).

In Uganda, the growth of Arabica coffee is favoured by temperatures which range between 18-21°C as compared to 22-30°C for Robusta coffee (Musoli et al., 2001). Arabica coffee is therefore more heat sensitive (Bunn, 2015) and thus will be more vulnerable to climate change (Jaramillo et al., 2009; Läderach et al., 2017). Therefore, the suitability area for growing this type of coffee will drastically reduce in future compared to Robusta coffee (Jaramillo et al., 2009; Jassogne et al., 2013).

¹ National Coffee Research Institute (NaCORI)/National Agricultural Research Organisation (NARO), P.O. Box 185 Mukono, Uganda. *Email : gkagezi@gmail.com

There will be a possible geographical shift by pests and diseases to higher altitudes due to global warming (Jaramillo et al., 2009), since elevation gradients can be used as analogues for global warming (Péré et al., 2013).

Rising temperatures have been reported to expand the altitudinal range of certain pests and diseases (Bongase, 2017). For example, the coffee berry borer (CBB), which has been having limited impact over 1500-1600 m.a.s.l in many countries (Baker and Hager, 2007; Jaramillo et al., 2009; Kyamanywa et al., 2012), has now been observed at altitudes more than 1800 m.a.s.l (Jaramillo et al., 2009; Kyamanywa et al., 2012; Agegnehu et al., 2015). Similarly, rising temperatures and rainfall have also been reported to have led to coffee leaf rust (CLR) becoming prevalent at higher altitudes than it used to be (Iscaro, 2014; Bebbber et al., 2016).

Basing on the above backdrop, we therefore conducted a study along an altitudinal gradient in Mt. Elgon, eastern Uganda in November-December 2016 to determine (i) the most prevalent insect pests and diseases influencing the production of Arabica coffee, (ii) the variation of insect pests and diseases along the altitudinal gradient, and, (iii) the insect pests and diseases that are likely to become a threat to Arabica coffee production under the future climate change scenarios. This information is a prerequisite for developing sustainable pest and disease control strategies as well as making decision for rational management practices.

2. Materials and methods

2.1 Description of the study area

A biological survey of pests and diseases was conducted in 4 districts located in the Mt. Elgon Arabica coffee growing region of Uganda, namely, Bulambuli, Kapchorwa, Sironko and Mbale in November-December 2016. This area is located between latitude 0.893° and 1.084°, and longitude 34.056° and 34.384° in the WGS84 coordinates system. The elevation ranges between 1084 and 2455 m.a.s.l (Jiang et al., 2014). The mean annual air temperature is about 23°C with average minimum and maximum temperatures of 15°C and 28°C, respectively. This region has a weak bi-modal rainfall pattern with a mean annual value of around 1500 mm (Bamutaze et al., 2010). The soils are generally deep and derived from volcanic ash as a product of a single weathering cycle (NEMA, 1998).

2.2 Sample selection and data collection

Ten (10) coffee growing households were purposively selected in each district and their coffee fields evaluated for pest and disease occurrence, severity and damage. Within each field, 30 coffee trees were selected along a diagonal. The first 10 coffee trees were assessed for both pests and diseases, then the other 20 assessed for only diseases. Pests were assessed on the coffee trees, leaves, clusters and berries. Trees were assessed for the presence or absence of white coffee stem borer (WCSB) – evidenced by either entry hole and/or the powdery frass resulting from burrowing of the larvae into the coffee stem, root mealy bugs – evidenced by either ant activity up and down the coffee plant and/or whitish powdery-like materials around the collar region and/or coffee roots as well as antestia bugs – by physical visual observation of the bugs.

To select the coffee leaves, clusters and berries to be sampled, the bearing head was divided into 3 imaginary sections – the upper, middle and lower. One primary branch with berries was randomly selected from each section. Then the total number of leaves as well as those damaged by the coffee leaf miner (CLM), skeletonizers, and lace bugs were determined. The total number of coffee clusters as well as those infested with canopy mealy bugs, canopy scales, and berry moth was determined on the sampled primary branch. The middle cluster of the sampled primary branch was selected and the number of berries as well as those infested with coffee berry borer (CBB) was determined. For the diseases, the incidence (presence or absence) and severity (scored on a scale of 1-5 where; 1=absent, 2= 1-10%, 3= 11-25%, 4= 26-50% and 5=>50% of leaves and/or berries are infected) were determined for coffee berry disease (CBD), coffee leaf rust (CLR), red blister disease and Elgon dieback.

2.3 Data analysis

Data for pest as well as disease incidence and severity were summarized using descriptive statistics including means and percentages. In addition, multiple regressions were run to assess the relationship between altitude and Arabica coffee pests and disease in order to determine those pests and diseases that are likely to be a threat under the future climate change scenarios. All the analyses were done in SAS v. 9.1 for Windows (SAS, 2008).

3. Results and Discussion

This section presents the results obtained in the study as well as providing the discussion on the observed results.

3.1 Insect pests recorded on Arabica coffee in Mt. Elgon region

A total of seven insect species, namely, the white coffee stem borers (WCSB), root and canopy mealy bugs, coffee leaf miners (CLM), leaf skeletonizers, lace bugs and coffee berry borers (CBB) were recorded on Arabica coffee in the Mt. Elgon region of Uganda (Table 1). This finding corroborates with several studies in this region (Kyamanywa et al., 2006; Egonyu et al., 2015; Jonsson et al., 2015; Kucel et al., 2016; Liebig et al., 2016).

Table 1: Percentage of trees, leaves, cluster and berries infested by the various Arabica coffee pests in Mt. Elgon region of Uganda

District	Infested trees (%)		Infested leaves (%)			Infested clusters (%)	Infested berries (%)
	WCSB	RMB	LB	SKL	CLM	CMB	CBB
Bulambuli	27.3	16.4	5.9	3.5	0.4	2.8	0.7
Kapchorwa	0.0	0.0	2.2	0.0	0.7	0.0	1.6
Mbale	13.0	19.0	7.8	3.6	0.3	0.5	1.9
Sironko	10.0	7.0	2.0	6.5	0.3	0.3	2.1
Overall	12.6	10.6	4.5	3.5	0.4	1.0	1.5

WSB=White coffee stem borers, RMB=Root mealy bugs, LB=Lace bugs, SKL=Skeletonizers, CLM=Coffee leaf miners, CMB=Canopy mealy bugs, CBB=Coffee berry borer

WCSB was the most prominent insect pest recorded, scoring the highest incidence of 13%. The importance of this insect pest in this region has been reported in biological surveys (Kyamanywa et al., 2006; Egonyu et al., 2015; Kucel et al., 2016) as well as farmers' perceptions (Liebig et al., 2016). WCSB has also been observed to be a major problem to Arabica coffee production in the Kilimanjaro region of Tanzania (Magina et al., 2010) as well as in several Asian countries (Venkatesha and Dinesh, 2012). Damage by this pest species is caused by its larva boring into the stem of coffee, weakening it and leading to yellowing of the foliage (Le Pelley, 1968). Cumulative yield losses of more than 25% in coffee due to this pest have been reported in Zimbabwe (Kutywayo et al., 2013).

3.2 Diseases recorded on Arabica coffee in Mt. Elgon region

On the other hand, three diseases, namely: - brown eye spot (BES), coffee leaf rust (CLR) and coffee berry disease (CBD) were observed in the region (Table 2). These diseases have been reported to be dominating the disease spectrum in the region (UCDA, 2011; Matovu et al., 2013).

Table 2: Severity (on a 1-5 scale) of Arabica coffee diseases in the Mt. Elgon region, eastern Uganda

	Coffee leaf rust (CLR)	Brown eye spot disease (BES)	Coffee Berry Disease (CBD)
Bulambuli	1.8	1.5	1.2
Kapchorwa	1.7	1.4	1.8
Mbale	1.8	1.7	1.0
Sironko	2.2	1.9	1.0
Overall	1.9	1.6	1.2

CLR registered the highest severity scores of 1.9 on a 1-5 scale. The importance of this disease in Mt. Elgon region has been recognized by biological studies by Matovu et al. (2013) as well as farmers' perceptions (Luzinda et al., 2015; Liebig et al., 2016). Similarly, CLR is one of the most important diseases of coffee worldwide (Hindorf and Omondi, 2011). The disease is caused by the fungus, *Hemileia vastatrix* (Berkeley and Broome) and manifests itself as yellow pustules on the lower surface of leaves, turning them to orange-yellow. It causes defoliation, leading to loss of yield of 10-50% and quality of coffee (Kushalappa and Eskes, 1989; van der Vossen, 2001; Silva et al., 2006).

3.3 Variation of the incidence of Arabica coffee insect pests with altitude

Results showed that, altitude significantly ($p < 0.05$) influenced the incidence of only WCSB and CLM among the insect pest species (Table 3). The incidence of WCSB decreased significantly ($p = 0.0169$) with increasing altitude, a similar trend observed by Jonsson et al. (2015) in the Mt. Elgon area and elsewhere (Le Pelley, 1968; Wrigley, 1988; Waller et al., 2007).

Lawton et al. (1987) suggested four possible causes for this decrease in pest incidence and damage - reduced habitat area at high elevations, reduced resource diversity, increasingly unfavourable environments, and/or reduced primary productivity at higher altitudes.

Table 3: Relationship between Arabica coffee insect pests and altitude in Mt. Elgon region, eastern Uganda

Variable	Df	Parameter estimate	Standard error	t value	Pr> t
Intercept	1	1620.35424	68.81287	23.55	<.0001
White coffee stem borers (WCSB)	1	-4.7767	1.89169	-2.53	0.0169
Root mealy bugs (RMB)	1	1.06663	2.22415	0.48	0.6349
Coffee leaf miners (CLM)	1	123.37257	48.02125	2.57	0.0152
Leaf skeletonizers (LS)	1	-15.01215	9.02984	-1.66	0.1065
Lace bugs (LB)	1	-3.68124	6.53014	-0.56	0.5770
Canopy mealy bugs (CMB)	1	13.16578	10.39109	1.27	0.2146
Coffee berry borer (CBB)	1	-5.21794	29.90727	-0.17	0.8626

On the other hand, the incidence of CLM increased significantly ($p=0.0152$) with increasing altitude. Our finding agrees with studies by Lomeli-Flores et al. (2009) conducted in Mexico, but contradict results by Nestel et al. (1994) and Tuelher et al. (2003). The difference in CLM incidence and damage at different elevations may be explained in part by the variation of rainfall and ambient temperatures which affect the coffee plant and thus, the pest and its natural enemies (Nestel et al., 1994; Pereira et al., 2007; Lomeli-Flores et al., 2009).

3.4 Variation of the severity of Arabica coffee diseases with altitude

Table 4 shows that among the Arabica coffee diseases, the severity of only CBD varied significantly ($p=0.0004$) with increasing altitude (Table 4). This observation agrees with findings by Mulinge (1971), Zeru et al. (2009) and Matovu et al. (2013). This relationship could be due to varying temperatures and moisture conditions as altitude increases (Zeru et al., 2009; Hindorf and Omondi, 2011). This subsequently translates into successful spore dispersal, germination and colonization of the disease (Griffiths and Waller, 1971; Ntahimpera et al., 1999; MouenBedimo et al., 2012).

Table 4: Relationship between severity of Arabica coffee diseases and altitude in Mt. Elgon region, eastern Uganda

Variable	Df	Parameter estimate	Standard error	t value	Pr> t
Intercept	1	1677.59446	216.95591	7.73	<.0001
Brown eye spot (BES)	1	-240.81709	213.95077	-1.13	0.2678
Coffee leaf rust (CLR)	1	-82.42293	163.21746	-0.50	0.6166
Coffee berry disease (CBD)	1	374.99216	96.26465	3.90	0.0004

3.5 Insect pests and diseases that are likely to become a threat under the future climate change scenario

Our results showed that only the incidence of WCSB decreased significantly ($p=0.0169$) with increasing altitude (Table 3). Our finding is consistent with biological studies by Kyamanywa et al. (2006) and Jonsson et al. (2015) as well as farmers' perceptions by Liebig et al. (2016) in this region. Similarly, Kutwayo et al. (2013) observed that WCSB was more prevalent at low altitude and significantly increased with altitude in Zimbabwe. This implies that basing on the future climate change scenarios (Jassogne et al., 2013), this insect pest which is currently a problem at low altitudes is expected to extend its niche to higher altitudes with increase in temperatures (Jaramillo et al., 2009, 2011; Kutwayo et al., 2013; Ovalle-Rivera et al., 2015), since elevation gradients can be used as analogues for global warming (Péret et al., 2013).

Given this general increased risk from WCSB in the face of climate change, there is a need to develop adaptation strategies for minimizing its impact to the coffee crop. This should go hand in hand with the various stakeholders who derive their livelihood from the crop along the value chain (Jassogne et al., 2013; Kutwayo et al., 2013). One of the proven strategies for mitigating these impacts is the introduction of shade systems in the coffee gardens (Jaramillo et al., 2009, 2011; Morris et al., 2016; Vezy et al., 2016; Alemu and Dufera, 2017). Shade systems can reduce the temperatures in coffee gardens by 2-6°C (Barradas and Fanjul, 1986) and solar radiation (Lin, 2007), thus, buffering the coffee plants from microclimate variability (Beer et al., 1998).

However, these tree species should offer limited competition to the coffee plants as well as not promoting pests and diseases (Beer et al., 1998; Kagezi et al., 2015; Netsere and Kufa, 2015; Likassa and Gure, 2017).

4. Conclusion

A total of seven insect pest species were observed on Arabica coffee in the Mt. Elgon region of Uganda - with the white coffee stem borers having the highest incidence of 13%. On the other hand, three diseases were observed – with CLR having the highest severity of 1.9. The incidence of WCSB and CLM as well as the severity of CBD varied significantly ($p < 0.05$) with increasing altitude. However, it was only the incidence of WCSB that decreased significantly ($p = 0.0169$) with increasing altitude, implying that this insect pest which is currently a major pest at low altitude will become a threat at higher altitudes under the future climate scenarios. Therefore, adaptation mechanisms for climate adversaries such as planting shade trees in coffee should be developed and promoted. However, these shade systems should offer limited competition to the coffee plant as well as not promote pests and diseases.

Acknowledgement

This study was funded by the Government of Uganda (GoU) through the Uganda Coffee Development Authority (UCDA). We also acknowledge the contribution of Robert Ekwaru, Viola Kigonya, Ruth Nseranyi and Julie Nabirye during data collection as well as the farmers participated in the study.

References

- Agegnehu, E., Thakur, A. & Mulualem, T. (2015). Potential impact of climate change on dynamics of coffee berry borer (*Hypothenemus hampei* Ferrari) in Ethiopia. Open Access Library, 2, e1127.
- Alemu, A. & Dufera, E. (2017). Climate smart coffee (*Coffea arabica*) production. American Journal of Data Mining and Knowledge Discovery, 2(2), 62-68.
- Baker, P. & Hagar, J. 2007. Global warming: Effects on global coffee. Long Beach, SCAA Conference Handout.
- Bamutaze, Y., Tenywa, M.M., Majaliwa, M.J.G., Vanacker, V., Bagoora, F., Magunda, M., Obando, J. & Wasige, E.E. (2010). Infiltration characteristics of volcanic sloping soils on Mt. Elgon, Eastern Uganda. Catena, 80, 122-130.
- Barradas, V.L. & Fanjul, L. (1986). Microclimatic characterization of shaded and open-grown coffee (*Coffea arabica* L.) plantations in Mexico. Agricultural and Forest Meteorology, 38, 101-112.
- Bebber, D.P., Castillo, A.D. & Gurr, S.J. (2016). Modelling coffee leaf rust risk in Colombia with climate reanalysis data. Philosophical Transactions of the Royal Society B, 371, 20150458.
- Beer, J., Muschler, R., Kass, D. & Somarriba E. (1998). Shade management in coffee and cacao plantations. Agroforestry Systems, 38, 139-164.
- Bongase, E.D. (2017). Impacts of climate change on global coffee production industry: Review. African Journal of Agricultural Research, 12(19), 1607-1611.
- Bunn, C. (2015). Modeling the climate change impacts on global coffee production. PhD thesis, Humboldt-Universität zu Berlin.
- Deutsch, C.A., Tewksbury, J.J., Huey, R.B., Sheldon, K.S., Ghalambor, C.K., Haak, D.C. & Martin, P.R. (2008). Impacts of climate warming on terrestrial ectotherms across latitude. PNAS, 105(18), 6668-6672.
- Egonyu, J.P., Kucel, P., Kagezi, G.H., Kovach, J., Rwomushana, I., Erbaugh, M., Wekono, Salifu, D. & Kyamanywa, S. (2015). *Coffea arabica* variety KP423 is less susceptible to the cerambycid coffee stem-borer *Monochamus leuconotus*, but common stem treatments seem ineffective against the pest. African Entomology, 23(1), 68-74.
- Griffiths, E. & Waller, J.M. (1971). Rainfall and cropping patterns in relation to coffee berry. Annals of Applied Biology, 67, 75-91.
- Hill, R.V. (2005). Risk, production and poverty: A study of coffee in Uganda. D. Phil thesis, University of Oxford.
- Hindorf, H. & Omondi, C.O. (2011). A review of three major fungal diseases of *Coffea arabica* L. in the rainforests of Ethiopia and progress in breeding for resistance in Kenya. Journal of Advanced Research, 2, 109-120.
- Hodkinson, I.D. (2005). Terrestrial insects along elevation gradients: species and community responses to altitude. Biological Reviews, 80, 489-513.
- Iscaro, J. (2014). The Impact of climate change on coffee production in Colombia and Ethiopia. Global Majority E-Journal, 5(1), 33-43.

- Jaramillo, J., Chabi-Olaye, A., Kamonjo, C., Jaramillo, A., Vega, F.E., Poehling, H.M. & Borgemeister, C. (2009). Thermal tolerance of the coffee berry borer *Hypothenemus hampei*: predictions of climate change impact on a tropical insect pest. *PLoS One*, 4(8), e6487.
- Jaramillo, J., Muchugu, E., Vega, F.E., Davis, A., Borgemeister, C. & Chabi-Olaye, A. (2011). Some like it hot: The influence and implications of climate change on coffee berry borer (*Hypothenemus hampei*) and coffee production in East Africa. *PLoS One*, 6(9), e24528.
- Jassogne, L., Läderach, P. & van Asten, P. (2013). The Impact of Climate Change on Coffee in Uganda: Lessons from a case study in the Rwenzori Mountains. *Oxfam Policy and Practice: Climate Change and Resilience*, 9, 51-66.
- Jiang, B., Bamutaze, Y. & Pilesjö, P. (2014). Climate change and land degradation in Africa: a case study in the Mount Elgon region, Uganda. *Geo-spatial Information Science*, 17(1), 39-53.
- Jonsson, M., Raphael, I.A., Ekbom, B., Kyamanywa, S. & Karungi, J. (2015). Contrasting effects of shade level and altitude on two important coffee pests. *Journal of Pest Science*, 88, 281-287.
- Kagezi, G., Kucel, P., Egonyu, J.P., Kyamanywa, S., Karungi, J.T., Pinard, F., Jaramillo, J., van Asten, P.J.A., Wagoire, W.W. & Ngabirano, H. (2015). A review of the status and progress in management research of the black coffee twig borer, *Xylosandrus compactus* (Eichhoff) in Uganda. In: *Proceeding of the 25th International Conference on Coffee Science*. ASIC. Paris: ASIC, pp. 42-52.
- Kucel, P., Kagezi, G.H., Rwomushana, I., Kyamanywa, S., Kovach, J., Roberts, A. & Erbaugh, M. (2016). Distribution and status of Arabica coffee pests in the Mountain Elgon area of Uganda. In: *Proceedings of the 1st NARO-MAK Joint Agricultural Dissemination Conference*, Munyonyo, Uganda.
- Kushalappa, A.C. & Eskes, A.B. (1989). *Coffee rust, epidemiology and control*. CRC Press, Inc. Boca Raton, Florida, USA.
- Kutywayo, D., Chemura, A., Kusena, W., Chidoko, P. & Mahoya, C. (2013). The impact of climate change on the potential distribution of agricultural pests: The case of the coffee white stem borer (*Monochamus leuconotus* P.) in Zimbabwe. *PLoS One*, 8(8), e73432.
- Kyamanywa, S., Ogwang, J.A. & Kucel, P. (2006). Biological monitoring of Arabica coffee insect pests in the Mount Elgon area. Report 10 pp.
- Kyamanywa, S., Kucel, P., Kagezi, G., Nafuna, K., Ssemwogerere, C., Kovach, J. & Erbaugh, M. (2012). IPM of the white stem borer and root mealy bugs on Arabica coffee in the Mt. Elgon region in Uganda. In: *Proceedings of the 7th International Integrated Pest Management Symposium IPM on the World Stage*, Memphis, Tennessee.
- Laderach, P., Ramirez-Villegas, J., Navarro-Racines, C., Zelaya, C., Martinez-Valle, A., & Jarvis, A. 2017. Climate change adaptation of coffee production in space and time. *Climatic Change*, 141(1), 47-62.
- Lawton, J.H., Macgarvin, M. & Heads, P.A. (1987). Effects of altitude on the abundance and species richness of insect herbivores on bracken. *Journal of Animal Ecology*, 56, 147-160.
- Le Pelley, R.H. (1968). *Pests of coffee*. Longmans, Green and Co., London.
- Liebig, T., Jassogne, L., Rahn, E., Läderach, P., Poehling, H-M., Kucel, P., van Asten, P. & Avelino, J. (2016). Towards a collaborative research: A case study on linking science to farmers' perceptions and knowledge on Arabica coffee pests and diseases and its management. *PLoS One*, 11(8), e0159392.
- Lin, B.B. (2007). Agroforestry management as an adaptive strategy against potential microclimate extremes in coffee agriculture. *Agricultural and Forest Meteorology*, 144, 85-94.
- Lomeli-Flores, J.R., Barrera, J.F. & Bernal, J.S. (2009). Impact of natural enemies on coffee leaf miner *Leucoptera coffeella* (Lepidoptera: Lyonetiidae) population dynamics in Chiapas, Mexico. *Biological Control*, 51, 51e60.
- Luzinda, H., Nelima, M., Wabomba, A., Kangire, A., Musoli, P.C. & Musebe, R. (2015). Farmer awareness, coping mechanisms and economic implications of coffee leaf rust disease in Uganda. *Uganda Journal of Agricultural Science*, 16(2), 207-217.
- Magina, F.L., Makundi, R.H., Maerere, A.P., Maro, G.P. & Teri, J.M. (2010). Temporal variations in the abundance of three important insect pests of coffee in Kilimanjaro region, Tanzania. In: *23rd International Conference on Coffee Science*. Bali, Indonesia. October 3-8, 2010. ASIC, Paris.
- Matovu, R., Kangire, A., Phiri, N.A., Hakiza, G.J., Kagezi, G.H. & Musoli, P.C. (2013). Ecological factors influencing incidence and severity of coffee leaf rust and coffee berry disease in major Arabica coffee growing districts of Uganda. *Uganda Journal of Agricultural Sciences*, 14(1), 87-100.
- Morris, K.S., Méndez, V.E. van Zonneveld, M., Gerlicz, A. & Caswell, M. 2016. Agroecology and climate change resilience in smallholder coffee agroecosystems of Central America. *ARLG Research Brief # 4*. Agroecology and Rural Livelihoods Group (ARLG), University of Vermont: Burlington, VT.

- Mouen Bedimo, J.A., Dufour, B.P., Cilas, C. & Avelino, J. (2012). Effects of shade trees on Coffee Arabica pests and diseases. *Cahiers Agricultures*, 21, 89-97.
- Mulinge, S.K. (1971). Effect of altitude on the distribution of the fungus causing coffee berry disease in Kenya. *Annals of Applied Biology*, 67, 93-98.
- Musoli, P.C., Hakiza, G.J., Birinkunzira, J.B., Kibirige-Sebunya & Kucel, P. (2001). Coffee (*Coffea* spp.). In: Agriculture in Uganda Vol. II. Mukiibi, J.K. (ed.). Fountain Publishers/CTA/NARO. Kampala, Uganda. pp 376-436.
- NEMA (1998). National Environmental Management Authority (NEMA) State of the Environment Report for Uganda.
- Nestel, D., Dickschen, F. & Altieri, M.A. (1994). Seasonal and spatial population loads of a tropical insect: the case of the coffee leaf miner in Mexico. *Ecological Entomology*, (19), 159- 167.
- Netsere, A. & Kufa, T. (2015). Determining suitable shade trees, planting pattern and spacing for Arabica coffee production in south and south-western Ethiopia. *Journal of Biology, Agriculture and Healthcare*, 5(17), 9-15.
- Ntahimpera, N., Wilson, L.L., Ellis, M.A. & Madden, L.V. (1999). Comparison of rain effects on splash dispersal of three *Colletotrichum* species infecting strawberry. *Phytopathology*, 89, 555-563.
- Ohmura, A. (2012). Enhanced temperature variability in high-altitude climate change. *Theoretical and Applied Climatology*, 110, 499-508.
- Ovalle-Rivera, O., Läderach, P., Bunn, C., Obersteiner, M. & Schroth, G. (2015). Projected shifts in *Coffea arabica* suitability among major global producing regions due to climate change. *Plos One*, 10(4), e0124155.
- Péré, C., Jactel, H. & Kenis, M. (2013). Response of insect parasitism to elevation depends on host and parasitoid life-history strategies. *Biology Letters*, 9(4), 20130288, 4 pp.
- Pereira, E.J., Picanço, M.C., Bacci, L., Crespo, A.L. & Guedes, R.N.C. (2007). Seasonal mortality factors of the coffee leaf miner, *Leucoptera coffeella*. *Bulletin of Entomological Research*, 97, 421e432.
- SAS (2008). SAS/STAT Software: Version 9.2, Cary, NC: SAS Institute Inc.
- Silva, M.C., Varzea, V., Guimaraes, L.G., Azinheira, H.G., Fernandez, D., Petitot, A.S., Lashermes, P., Nicole, M. & Bertrand, B. (2006). Coffee resistance to the main diseases: Leaf rust and coffee berry disease. *Brazilian Journal of Plant Physiology*, 18(1), 119-147.
- UCDA (2011). Uganda Coffee Development Authority (UCDA). UCDA September 2011 monthly report.
- van der Vossen, H.A.M. (2001). Coffee breeding practices. In: R.J. Clarke & O.G. Vitzthum (Eds.) *Coffee recent developments*. Blackwell Science, Oxford, pp. 184-201.
- Venkatesha, M.G. & Dinesh, A.S. (2012). The coffee white stem borer *Xylotrechus quadripes* (Coleoptera: Cerambycidae): bioecology, status and management. *International Journal of Tropical Insect Science*, (4).177–188.
- Vezy, R., Picart, D., Christina, M., Soma, M., Georgiou, S., Charbonnier, F., Loustau, D., Imbach, P.B., Hidalgo, H.G., Alfaro, E.J., le Maire, G. & Rouspard, O. (2016). Effect of shade on temperature mitigation and canopy assimilation of coffee agroforestry systems. In: *Proceedings of the 3rd European Agroforestry Conference Montpellier SupAgro, France*.
- Waller, J.M, Bigger, M. & Hillocks, R.J. (2007). *Coffee pests, diseases and their management*. Oxfordshire UK. CAB International.
- Wrigley, G. (1988). *Coffee*. Tropical Agriculture Series. Longman, London.
- Zeru, A., Assefa, F., Aduagna, G. & Hindorf, H. (2009). Occurrence of fungal diseases of *Coffea arabica* L. in montane rain forests of Ethiopia. *Journal of Applied Botany and Food Quality*, 82(2), 148-151.